Application No. 10/353,059

Amendment dated: September 25, 2008

Reply to Non-Final Office Action of May 30, 2008

AMENDMENTS TO THE CLAIMS

This Listing of Claims replaces all prior versions and listings of claims in the application.

- 1. (currently amended) A crystal <u>form</u> of 6-fluoro-1-methyl-7-[4-(5-methyl-2-oxo-1,3-dioxolen-4-yl)-methyl-1-piperazinyl]-4-oxo-4H-[1,3]thiazeto[3,2-a]quinoline-3-carboxylic acid acetonitrile solvate, which has diffraction peaks at least at 7.3, 14.7, 19.2 and 22.3 in the powder X-ray diffraction spectrum.
 - 2. (canceled)
- 3. (currently amended) A method for producing a crystal <u>form</u> of 6-fluoro-1-methyl-7-[4-(5-methyl-2-oxo-1,3-dioxolen-4-yl)-methyl-1-piperazinyl]-4-oxo-4H-[1,3]thiazeto[3,2-a]quinoline-3-carboxylic acid acetonitrile solvate, characterized by which comprises performing crystallization from an acetonitrile solution of 6-fluoro-1-methyl-7-[4-(5-methyl-2-oxo-1,3-dioxolen-4-yl)-methyl-1-piperazinyl]-4-oxo-4H-[1,3]thiazeto[3,2-a]quinoline-3-carboxylic acid by controlling its supersaturation concentration (g/100 g) to be from 2.15 to 2.36 at the time of occurrence of spontaneous nucleus nucleation.
- 4. (currently amended) A method for producing a crystal <u>form</u> of 6-fluoro-1-methyl-7-[4-(5-methyl-2-oxo-1,3-dioxolen-4-yl)-methyl-1-piperazinyl]-4-oxo-4H-[1,3]thiazeto[3,2-a]quinoline-3-carboxylic acid acetonitrile solvate, which comprises performing crystallization from an acetonitrile solution of 6-fluoro-1-methyl-7-[4-(5-methyl-2-oxo-1,3-dioxolen-4-yl)-methyl-1-piperazinyl]-4-oxo-4H-[1,3]thiazeto[3,2-a]quinoline-3-carboxylic acid by controlling its supersaturation concentration (g/100 g) to be from 0.41 to 2.36 at the time of addition of a seed crystal.
- 5. (Original) The method according to Claim 4, wherein the solution at the time of addition of the seed crystal has a temperature of 70°C or lower.

Application No. 10/555,039 Docket No.: 03702/0203547-US0

Amendment dated: September 25, 2008

Reply to Non-Final Office Action of May 30, 2008

6. (currently amended) A method for producing the type III crystal <u>form</u> of 6-fluoro-1-methyl-7-[4-(5-methyl-2-oxo-1,3-dioxolen-4-yl)-methyl-1-piperazinyl]-4-oxo-4H-[1,3]thiazeto[3,2-a]quinoline-3-carboxylic acid, which comprises desolvating a crystal of 6-fluoro-1-methyl-7-[4-(5-methyl-2-oxo-1,3-dioxolen-4-yl)-methyl-1-piperazinyl]-4-oxo-4H-[1,3]thiazeto[3,2-a]quinoline-3-carboxylic acid acetonitrile solvate.